

# cDNA Synthesis Kit

Manual LS62 Version 1.2 April 2017

Store at -20°C



### Product description

Reverse transcription

The ability to reverse the process of translation from DNA to RNA has been a crucial application of modern molecular biology. The enzyme, a RNA-dependent DNA-Polymerase is called reverse transcriptase. Reversing the messenger RNA delivers the complementary DNA, which can be detected or quantified using PCR and qPCR approaches.

The lack of the  $3' \rightarrow 5'$  exonuclease activity made the MuLV increasingly the enzyme of choice. The MuLV has also been used as a template to engineer new enzymes with additional features. The FastGene Scriptase Basic is an enhanced version of the original MuLV RT.

### Kit Content

(100 Reactions)

- FastGene®Scriptase Basic (200 U/μl)
- 10x FastGene® Scriptase Basic buffer
- dNTP Mixture (2 mM each)
- Sterile water (RNase free)
- RNase Inhibitor(40 U/µl)
- Oligo dT (80 μM)
- Random Hexamer(100 µM)

#### **Quality Control**

- Purity: >99% on SDS-PAGE
- Endonuclease-free
- Exonuclease-free
- RNase-free
- · Inhibitor-free
- Satisfactory Yield

#### **Protocol**

#### Concentration of template RNA

- Total RNA: 1 ng-5 μg
- Messenger RNA (mRNA): 1 ng-0.25 μg
- Specific RNA: 0.01 pg-0.5 μg
- Mix the recommended concentration of RNA template with 1 μl oligo dT primer (80 μM) and/or 1 μl random hexamer (100μM)
- 2. Incubate at 42 °C for 5 minutes.

Note: Add an additional annealing step of 10 minutes at 25°C if random hexamers are used.

3. Store the RNA mixture on ice and add followed components:

10x FastGene® Scriptase Basic buffer		2 μΙ
FastGene® Scriptase Basic (200 units/μl)		
dNTP mixture (2 mM each)		2 μΙ
RNA-Template		
Primer		1-2 µl
RNase Inhibitor		
Distilled Water	up to	20 μΙ

4. Incubate at 42°C for 60 min.

<u>Note:</u> FastGene $^*$  Scriptase Basic can perform optionally over the full range of 42°C-50°C after optimization.

5. Incubate 90 °C for 5 min for complete enzyme deactivation.



## **cDNA Synthesis**

Please contact us for additional information:

#### info@nippongenetics.eu

Please contact our technical support:

#### support@nippongenetics.eu

LS52	FastGene® Scriptase Basic (20.000 units at 200 U/μl)	100 Reactions
LS62	FastGene® Scriptase Basic cDNA Synthesis containing Oligo dTs, random hexamer and RNase inhibitor	100 Reactions
LS53	FastGene® Scriptase II (20.000 units at 200 U/μl)	100 Reactions
LS63	FastGene® Scriptase II cDNA Synthesis containing Oligo dTs, random hexamer and RNase inhibitor	100 Reactions

