

# MIDORI Green Advance TBE Agarose Tablets

Cat. No. AG09

## Description

MIDORI Green Advance TBE Agarose Tablet contains a pre-determined amount of standard melting point agarose, MIDORI Green Advance stain, and TBE powder.

Midori Green Advance DNA stain emits green fluorescence when bound to DNA or RNA. It is compatible with a wide variety of gel reading instruments.

The purity of the agarose leads to an excellent transparency and a low background, important to obtain sharp and well-defined DNA/RNA bands with the highest sensitivity in the low molecular weight range.

## Safety

Caution when using hot, viscous solutions! Use suitable safety gear and open bottle gently to avoid accidents.

Midori Green Advance DNA stain is non-carcinogenic and according to the Ames test it causes significantly fewer mutations than Ethidium bromide. It can irritate skin and eyes. Please wear gloves while handling. A detailed safety report can be downloaded at [www.nippongenetics.eu](http://www.nippongenetics.eu).

## Specifications

- Melting point:  $88 \pm 1,5$  °C
- Separation range: 100 bp to >30 kb
- Product size: 100 tablets (0.5 g each)

## Applications

- Ideal for routine DNA and RNA gel electrophoresis and blotting assays.
- Convenient tablet format - no complicated messy weighing required.
- Fast dissolving protocol.

## NIPPON Genetics EUROPE GmbH

+49 2421 554960    info@nippongenetics.eu  
+49 2421 55496 11    www.nippongenetics.eu

## Protocol

1. Use the bottle or flask that is at least 3 times the volume of the solution being prepared.
2. Add an appropriate number of agarose tablets in the **water**. See the table below to achieve the needed gel percentage.

Gel %	1 tablet	2 tablets	3 tablets
1%	25 ml H <sub>2</sub> O	50 ml H <sub>2</sub> O	75 ml H <sub>2</sub> O
1,5 %	16.5 ml H <sub>2</sub> O	33 ml H <sub>2</sub> O	49.5 ml H <sub>2</sub> O
2 %	12.5 ml H <sub>2</sub> O	25 ml H <sub>2</sub> O	37.5 ml H <sub>2</sub> O

- Soak the tablet in the pure cold water for at least 3 minutes (or until it is dissolved) before heating. For tablet dissolving use water which is at RT. DO NOT use hot water.
- Heat the solution until it is clear and visually all the particles are dissolved.
- Cool the gel to 60-70°C.
- Cast the gel into the gel tray. The thickness of the gel should be <0.5cm.
- Run the gel in TBE running buffer.
- Detect the bands under Blue/Green LED or UV illuminator.

## Storage

Storage and shipping at RT. Protected from light.

## Ordering information

Cat. No.	Pack Size
AG09	100 pcs
AG09s	4 pcs